

Chemical composition of Nigella sativa Linn : Part 2 Recent advances

KHAN, Akram <<http://orcid.org/0000-0002-0386-3667>> and AFZAL, M

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/12364/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

KHAN, Akram and AFZAL, M (2016). Chemical composition of Nigella sativa Linn : Part 2 Recent advances. *Inflammopharmacology*, 24 (2), 67-79.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Chemical Composition of *Nigella sativa* Linn: Part 2 Recent Advances

M. AKRAM KHAN^{*1}; M. AFZAL²

¹ Biomolecular Science Centre, Sheffield Hallam University, Howard Street, Sheffield, S1 1WB, UK

² Department of Biological Sciences, Faculty of Science, Kuwait University, Kuwait

Abstract– This review article is an update on the previous article published on *Nigella sativa* Linn in this journal in 1999. It covers the medicinal properties of *Nigella sativa* and chemical syntheses of the alkaloids isolated from the seeds of the herb.

1.0 INTRODUCTION

The chemical composition and biological properties of *Nigella sativa* L. have previously been reviewed [Khan, 1999; Paarakh, 2010; Ahmad A, et. al., 2013]. In the previous review (Khan, 1999) were reported the large variety of organic compounds that are present in the seeds of *Nigella sativa* L. The seeds of this herb are used in the Middle East and South Asian countries for the treatment of a large variety of ailments and are accepted as a panacea. For example, the seeds or oil from the seeds have been used to control diabetes, hypertension, cancer (leukeamia, liver, lung, kidney, prostate, breast, cervix, skin), inflammation, hepatic disorder, arthritis, kidney disorder, cardiovascular complications and dermatological conditions (Khan *et al.*, 2011; Khan *et al.*, 2003b). A GC-MS analysis of the seed extract has shown it to be a mixture of eight fatty acids and 32 volatile terpenes. The major terpenes, thymoquinone (TQ), dithymoquinone (DTQ), trans-anethol, p-cymene, limonine, and carvone have been identified (Nickavar *et al.*, 2003). TQ and DTQ are both cytotoxic for various types of tumors (Worthen *et al.*, 1998). In addition diterpenes, triterpene and terpene alkaloids have been identified in *Nigella sativa* seeds. The methanolic extract of the seeds contain two types of alkaloids whilst the major principal active ingredient isolated from the volatile oil of *Nigella sativa* L. is TQ. Since *Nigella sativa* L. acts as a panacea exhibiting a wide variety of pharmacological actions discussed previously and updated in this report, interest has arisen in the total synthesis of the alkaloids isolated having the isoquinoline and indazole motifs. The

*Correspondent author. E-mail: m.a.khan@shu.ac.uk

isoquinoline alkaloids include nigellicimine (1) and nigellicimine- N-oxide (2), and the indazole alkaloids include nigellidine (3) and nigellicine (4) (Fig. 1). Since the previous review several new dolabellane-type diterpene alkaloids, nigellamines A₁-A₅ (5) have also been isolated from the methanolic extract of the seeds of *Nigella sativa* L. which have also received synthetic interest (Fig. 1). In this update on *Nigella sativa* we want to discuss the chemistry of these various alkaloids and TQ under separate headings.




Figure 1. Structures of alkaloids isolated from *Nigella sativa* L.

1.1 Pyrazole and Indazole ring systems




Figure 2. Types of indazole ring compounds

Indazole and pyrazole motifs are embedded in numerous pharmaceuticals and agrochemicals with a broad range of biological activities such as (6) (Penning et al, 1997), (7) (Plosker and Goa, 1991), (8) (de Paulis et al, 2006), (9) (Okuno et al, 2004), (10) (Maxwell, 2000) and (11) (Lahm et al, 2009) shown in Figure 3.




Figure 3. Structures of some pharmaceuticals and agrochemicals with indazole and pyrazole motifs

As a result of these biological activities being associated with the presence of pyrazole and indazole pharmacore in therapeutic compounds, the two indazole alkaloids nigellidine (3) and nigellicine (4) have attracted the attention of synthetic organic chemists for their total syntheses. Thus multigram quantities of these two alkaloids can now be obtained via their total syntheses that should enable their individual therapeutic evaluation to be possible.

2.0 CHEMISTRY OF THE ALKALOIDS AND TQ IN *Nigella sativa*

2.1 Total synthesis of Nigellidine (3)

The development of an efficient synthetic method using Pd(II)/Phen catalyst and conditions for the direct C-3 C-H arylation of (1H) indazole and pyrazole with ArI or ArBr was applied to the synthesis of nigellidine as shown in scheme 1 (Ye et al, 2013). The THP derivative of the commercially available 4-methoxy-6-methyl-(1H)-indazole was reacted with 4-bromoanisole using the C-3 arylation reaction as a key step to form the adduct (14) in 54% isolated yield on the gram scale. Deprotection of the tetrahydropyranyl (THP) group gave (15) which N-alkylation with 1,4-dibromobutane to afford (16) that underwent intramolecular cyclised to furnish the precursor (17). Demethylation of (17) by treatment with BBr₃ afforded the natural product nigellidine (3) as the hydrobromide salt in an overall yield of 18%.

2.2 The Total syntheses of Nigellicine (4)


To date there have been two total syntheses of nigellicine reported. In the first synthesis shown in scheme 2 commercially available 2-chloro-5-methylphenol (17) was transformed into the protected amide (10) which on lithiation and acylation with diethyloxalate yielded the amide-ester (20) that cyclised on treatment with 6M HCl acid into the isatin (21) (Elliott et al, 2005). Protection of the keto group in isatin (21) as the dimethyl acetal (22) enabled direct amination at nitrogen to give the hydrazine derivative (22) which on treatment with aqueous acid rearranged into an indazole carboxylic acid that was esterified via an acid chloride into (23). Alkylation of indazole ester (23) with 1,4-dibromobutane and subsequent intramolecular cyclisation produced a 4-methoxy derivative of nigellicine which was deprotected with PBr₃ to give nigellicine in an overall yield of 18%.

The second total synthesis of nigellicine (4) shown in scheme 3 starts with commercially available 2,5-dimethylphenol (24) which was converted into 2-bromo-6-methoxy-4-methylbenzaldehyde (25) by Clive's method (Inamoto et al, 2007). Treatment of aldehyde (25) with KCN and ethyl chloroformate in the presence of benzyltrimethylammonium chloride (BTAC) and 18-crown-6 in a mixture of water and 1,2-dichloroethane produced an intermediate cyanohydrin carbonate ester which was subsequently converted to a-ketoester (26) by LiHMDS-induced rearrangement. The


reaction of (26) with p-toluenesulfonyl hydrazide gave the key intermediate hydrazone (27) as a mixture of *E*- and *Z*-isomers, which was separable by column chromatography to obtain the major trans isomer that was subsequently converted by Pd-catalysed cyclisation to the indazole (28). Alkylation of the deprotected compound (29) with 1,4-dibromobutane produced the intermediate (30) which underwent intramolecular cyclisation in hot ethanol to furnish the nigellicine ethyl ester hydrobromide salt (31). Finally treatment of ester (31) with PBr₃ caused cleavage of the ester group and deprotection of the methoxy group to give nigellicine (4).

2.3 Total synthesis of Nigellamine A₂ (5b)

The delabellane diterpenes are ubiquitous molecules that are produced by animals, plants, fungi and marine sources and have interesting array of biological activities. The alkaloids nigellamines A₁-A₅ which have been isolated from *Nigella sativa* L. belong to the delabellane family of diterpenes and show potent lipid metabolism-promoting activity (Morikawa et al, 2004a). These biologically active alkaloids have complex structural features and have attracted the attention of synthetic organic chemists for their total synthesis.




Scheme 1. Total synthesis of nigellidine as the hydrobromide salt



Scheme 2 Total synthesis of nigellicine

One enantioselective total synthesis of nigellamine A₂ has so far been reported (Bian et al, 2006). In this synthesis shown in scheme 4 the starting lactone-diene (32) was transformed in three steps and on a multigram scale into the allylic ester (33) as a key intermediate. Iodolactonisation of diene (33) produced (34) which on radical alkynylation furnished the propynyl lactone (35). Desilylation and reduction of (35) yielded the propynyl lactol (36) which upon *in situ* iodination and subsequent silylation afforded the vinyl iodide (37) in good yield. The remaining carbon atoms of the nigellamine skeleton were constructed through cross coupling with alkyl zinc reagent and a repeat methylalumination-iodination sequence of reactions to afford substrate (38). Oxidation with pyridinium chlorochromate (PCC) gave an aldehyde at position C₂ which upon sonification underwent Cr-mediated cyclisation with the vinyl iodide group at position




Scheme 3. Second synthesis of Nigellicine (4)

C₃ to generate the 11-membered compound (39). Reductive opening of the lactone and selective acylation of the primary alcohol gave the substrate (40). Oxidation of (40) with Shi's ketone catalyst and oxone proceeded region- and stereoselectively to produce the desired epoxide-diol as the major product which was acylated with nicotinic acid to furnish ent-nigellamine A₂ (5b).

2.4 Novel synthetic Thymoquinone analogues

The compound 5-isopropyl-2-methyl-1,4-benzoquinone is known as thymoquinone (TQ) (41) shown in scheme 5. TQ is the major active principle of the oil of *Nigella sativa* L. and has been shown to exhibit anti-tumor activity against breast, lung, prostate, liver, colon and pancreatic cancer. Thus interest has arisen to synthesise more potent analogues of TQ. Recently reported are the novel analogues of TQ consisting of compounds (44a-b) were synthesised in two steps from TQ Sodium azide added to TQ in acetic acid to afford the reduced product (42) which on reaction with the aldehydes


(43a-b) generated the Schiff bases (44a-b) (Yusufi et al, 2013). These analogues have shown superior proliferative activity, excellent chemo-sensitizing activity against pancreatic cancer *in vitro* and in combination with Gemcitabine.



Scheme 4. Total synthesis of nigellamine A₂

One serious drawback with TQ is its toxicity at high doses and poor water solubility which limit its usage as a therapeutic agent. In order to alleviate this problem various types of nanocarrier for thymoquinone have been synthesised (Ravindran et al, 2010; Ganea et al, 2010; Alam et al, 2012; Singh et al, 2013). One recent study has reported the synthesis of PAG coated NIPAAM nanoparticles that are encapsulated with TQ for direct hepato-targeting. NIPAAM is a thermosensitive nanopolymer which is widely used as a successful drug delivery system against various diseases and PAG is a galactosylated moiety that targets the liver by interacting with asialoglycoprotein receptor (ASGP-R)

present on the surface of hepatocytes and delivers the drug directly to the liver (Verma et al, 2013). The toxicity of the nanocarrier (NIPAAAM) at this concentration is almost negligible and due to the size of the nanoparticle being smaller than the already reported nanothymoquinone.



Scheme 5. Synthesis of Thymoquinone analogues as anticancer agents

This study clearly has demonstrated that the nanoparticles are able to carry bulk amounts of drug to the liver, and their direct targeting to ASGP-R receptors present on hepatocytes has resulted in significant hepatoprotection at a low dose level that is 1000 times lower than the naked TQ. This nanocarrier approach offers a promising prospect for the future against various liver diseases.

3.0 BIOLOGICAL ACTIVITIES OF *Nigella sativa*

3.1 The anti-inflammatory activities

In animal studies *Nigella sativa* shows dose-dependent suppression of nociceptive pain response and cestocidal activity. These activities are shown by TQ that acts through indirect activation of the supraspinal mu(1)- and kappa-opioid receptor subtypes (Abdel-Fattah et al., 2000; Akhtar et al., 1991). The antihypertensive principal TQ and other constituents of *Nigella sativa* are also protective agents against the chromosomal aberrations induced by schistosomiasis (Aboul-Ela, 2002; el el Tahir et al., 1993a).

These compounds are used in the control of arterial blood pressure, anticholinergic, antihistaminic, tracheal relaxation, control of asthma and in the treatment of other allergic diseases (Ahmed *et al.*, 2013; Al-Majed *et al.*, 2001; Boskabady *et al.*, 2004; Kalus *et al.*, 2003; Steinmann *et al.*, 1997).




Figure 4. Chemical structures of principal active ingredient isolated from the volatile oil of *Nigella sativa* L

Nigellone (dithymoquinone) is the carbonyl dimer of TQ present in *Nigella sativa* and it inhibits the release of histamine giving relief in asthmatic conditions (Chakravarty, 1993; el Tahir *et al.*, 1993b). The spasmolytic and bronchodilator activities of *Nigella sativa* are mediated possibly through calcium channel blockade (Gilani *et al.*, 2001). Physiologically important activities shown by *Nigella sativa* include analgesic, anti-inflammatory, antimicrobial, antifungal and antibacterial effects (Hanafy *et al.*, 1991; Khan *et al.*, 2003a; Morsi, 2000) and CNS activity of its aqueous extract and volatile oil components (Al-Ghamdi, 2001; Al-Naggar *et al.*, 2003; Hajhashemi *et al.*, 2004; Haq *et al.*, 1995). The neuroprotective activity of *Nigella sativa* on neurotransmitter leading to antiepileptic activity has also been described (Arafa *et al.*, 2013). TQ, through an opioid receptor-mediated, increases in GABAergic tone, exhibits anticonvulsant activity in the petit mal epilepsy (Hosseinzadeh *et al.*, 2004).

3.2 Antiulcer and anticancer properties

Ethanol induced ulcer in rats has been reduced by *Nigella sativa* extracts (El-Dakhakhny *et al.*, 2000a). Ischaemia/reperfusion are linked by free radical generation and this could be controlled by an administration of TQ which could offer gastroprotective effects against gastric lesions (El-Abhar *et al.*, 2003). The chemosensitising effect of TQ

in the treatment of 5-Fluorouracil induced gastric cancer has been reported (Lei *et al.*, 2012).

3.3 Hepato-protective antioxidant activities

The aqueous extract of *Nigella sativa* (NS) is hepato-protective against carbon tetrachloride induced oxidative hepatic damage suggesting powerful antioxidative properties of NS extract (Al-Ghamdi, 2003; el-Dakhakhny *et al.*, 2000a; Mansour *et al.*, 2001; Meral *et al.*, 2003). NS protects liver by inhibiting enzyme leakage from hepatocytes caused by toxic substances such as carbon tetrachloride (Kanter *et al.*, 2003a). Through its antioxidant action, TQ is known to inhibit 5-lipoxygenase and 5-hydroxy-eicosatetraenoic acid (5-HETE) products suggesting its use in inflammatory pathogenesis (El-Dakhakhny *et al.*, 2002a). Hyperhomocysteinemia (HHcy) has been linked with oxidative stress. Therefore, NS has been demonstrated to improve total antioxidant status in rats treated with methionine induced HHcy (El-Saleh *et al.*, 2004). The oxygen free radical generated by gentamicin pathogenesis causing hepatotoxicity and nephrotoxicity are quenched by oil and seeds of NS (Ali, 2004) and the ethanolic extracts of NS have the potential to protect against gamma-radiation induced oxidative damage (Rastogi *et al.*, 2010). It has been reported that TQ inhibits the leakage of hepatic enzymes and the intracellular depletion of GSH protecting liver (Daba *et al.*, 1998).

3.4 Immunomodulatory effect

NS has established immunosuppressive and cytotoxic properties (Islam *et al.*, 2004) and the pharmacological and therapeutical properties of NS have been reviewed by many workers (Ahmad *et al.*, 2013; Ali *et al.*, 2003; Swamy *et al.*, 2000). The splenocyte proliferation, macrophage function, and NK anti-tumor activity of NS have revealed the potent immunomodulatory properties of the *Nigella* seeds (Majdalawieh *et al.*, 2010). The NS seed oil also shows hepatoprotective action against hypervitaminosis A and humoral immune responses and non-specific cellular immune responses (Al-Suhaimi, 2012; Al-shatwi, 2014). The immune modulating effect of NS is mediated through direct

stimulation of macrophage phagocytic activity or lymphocytes activation (Fararh *et al.*, 2004; Haq *et al.*, 1999).

NS is a known immune stimulant that protects against many pathological conditions (Corder *et al.*, 2003; Fararh *et al.*, 2004). Thus, the powerful antioxidative and protective properties of TQ in proteinuria and hyperlipidemia associated with nephrotic syndrome have been evaluated (Badary *et al.*, 2000). Along with TQ and other terpenoid compounds such as carvacrol, trans-anethole and 4-terpineol with antioxidant properties have been reported for NS (Burits *et al.*, 2000). Carvacrol is a known inhibitor of human neutrophil elastase and may be useful agent in phytotherapy for injuries such as chronic obstructive pulmonary disease and emphysema (Kacem *et al.*, 2006). The volatile NS oil shows anti-oxytotic and sperm characteristics due to its antioxidant activities (Mansour *et al.*, 2013; Aqel *et al.*, 1996).

3.5 Effect on Blood sugar and lipid profile


Streptozotocin (STZ) treated animals respond to NS extracts with normalizing blood glucose through extrapancreatic actions rather than by stimulated insulin release and ascertain to be protective against type-2 diabetes (El-Dakhakhny *et al.*, 2002b; Fararh *et al.*, 2004; Fararh *et al.*, 2002; Hawsawi *et al.*, 2001). The significant increase in lipid peroxidation by STZ is also controlled by NS and has protective effect in diabetes by decreasing oxidative stress and regeneration/proliferation of the beta-cells in the islets of Langerhans (Kanter *et al.*, 2004; Kanter *et al.*, 2003b). A petroleum ether extract of NS exhibits insulin-sensitizing activity (Le *et al.*, 2004) and the mechanism of NS extract in the control of diabetes has been shown to be through controlled insulin release (Rchid *et al.*, 2004). At the same time, amendment in the blood lipids profile has been suggested by the use of NS extracts (El-Dakhakhny *et al.*, 2000a). Arachidonic acid induced blood platelet aggregation and blood coagulation are inhibited by NS indicating its potential use in thrombosis (Enomoto *et al.*, 2001). TQ is involved in the inhibition of arachidonic acid generated eicosanoids and lipid peroxidation (Houghton *et al.*, 1995).

3.6 Effect on arthritis

In human, TQ has been shown to be effective in rheumatoid arthritis (Gheita *et al.*, 2012). Inhibition of arachidonic acid generated eicosanoids (thromboxane B₂, leukotriene B₄) supports the use of NS for the treatment of rheumatoid arthritis and other inflammatory diseases (Houghton *et al.*, 1995). TQ has been implicated in bone healing in an animal model (Kirui *et al.*, 2004). Inhibition of leukotrienes through 5-lipoxygenase and LTC₄ synthase activities in eicosanoid pathway has been well documented (Mansour *et al.*, 2004).

3.7 Anticancer Activity of TQ

A number of antitumor compounds have been identified from NS. These compounds are TQ, alpha-hederin a triterpene, isopropylmethylphenols and dollabelane-type diterpene alkaloid nigellamine A3, A4, A5 and C (Kumara *et al.*, 2001; Michelitsch *et al.*, 2004; Morikawa *et al.*, 2004a; Morikawa *et al.*, 2004b). Thus, numerous types of cancers such as Ehrlich ascites carcinoma (EAC), Dalton's lymphoma ascites (DLA) and Sarcoma-180 (S-180) cells, colon carcinoma, pancreatic carcinoma and hepatic carcinoma have been treated with NS extracts *in vitro* (Salomi *et al.*, 1992; Samarakoon *et al.*, 2010). Changes in intracellular GSH and redox status for mitochondrial function are important factors in the mechanism of alpha-hederin induced cell death (Swamy *et al.*, 2003).



The NS extract exerts anti-hepatocarcinoma effect through modulation of apoptosis (Samarakoon *et al.*, 2012). The regulation of pro- and anti apoptic genes by NS has been demonstrated in treating cervical cancer (Shafi *et al.*, 2009). In many cases the antitumor activity of NS seeds has been attributed to the volatile component thymoquinone (structurally related to tert-butylhydroquinone, a potent antioxidant) that has the potential to protect rat liver against diethylnitrosamine (DEN) induced hepatocarcinogenesis (Iddamaldeniya *et al.*, 2003). It also improves the therapeutic efficacy of ifosfamide by decreasing nephrotoxicity and improving antitumor activity (Badary, 1999; Saleem *et al.*, 2012).

TQ also affects the benzo-a-pyrene induced clastogenic activity in rats and 20-methylcycloanthrene induced fibrocarcinoma is inhibited by TQ present in NS extracts (Badary *et al.*, 2007; Badary *et al.*, 2001). While supplementation by NS and honey in the treatment of methylnitrosourea induced inflammation, carcinogenesis and oxidative stress has been reported (Mabrouk *et al.*, 2002), the lipid peroxidation induced liver damage in diabetic rats has also been mentioned (Meral *et al.*, 2001).

The pro-oxidant nitric oxide production is inhibited by NS extracts validating the fact that NS has anti-inflammatory activities (Mahmood *et al.*, 2003). Model *in vivo* experiments with Schistosomiasis mansoni infected mice have concluded that NS extract have a great protective potential against oxidative stress protecting liver (Mahmoud *et al.*, 2002). The mode of action of TQ against cancer has been suggested to be through its antioxidative properties and interaction with DNA synthesis. The antioxidant and pro-oxidant properties of TQ have been substantiated by augmented TQ mediated scavenging of superoxide anion (Badary *et al.*, 2003). However, presence of the phenolic compounds in NS, such as vanillic acid, could also contribute to the antioxidant properties of NS. These compounds may also be responsible for its antimutagenic activities (Bourgou *et al.*, 2008; Khader *et al.*, 2010). TQ exhibits advanced antimyeloma activity in MDN and XD2 multiple myeloma malignant plasma cells (Badr *et al.*, 2011). However, the mechanism of chemotaxis of malignant plasma cells is not well defined.

3.8 Effect of TQ on Pancreatic carcinoma (PC)

PC is one of the most deadly cancers with almost invariably fatal consequences. TQ has antitumor activity against PC. To combat PC, the dose of TQ has to be high. Therefore, many attempts have been made to study structure activity relationships by synthesizing TQ analogs and some of these compounds have potent antitumor activity against PC (Banerjee *et al.*, 2010). Gemcitabine- or oxaliplatin-induced activation of NF-kappaB is inhibited by TQ, resulting in the chemosensitization of pancreatic tumors to conventional therapeutics (Banerjee *et al.*, 2009). Progressive apoptosis is also inhibited by NS (Corder *et al.*, 2003).

3.9 Co-administration of NS with other substances

Cisplatin is a widely used drug that induces kidney toxicity. It has been established that when cisplatin is co-administered with NS, the nephrotoxicity is reduced (El-Daly, 1998; Nair *et al.*, 1991; Ulu *et al.*, 2012). A co-administration of NS with green tea extract prevents cytotoxicity of organophosphorus compounds (Korany *et al.*, 2011). Co-administration of saffron with NS in the treatment of chemical carcinogenesis has also been reported (Salomi *et al.*, 1991).

4.0 Breast Cancer

It is one of the most common causes of death in women and there is no effective treatment except mastectomy. Many substances have been shown to have mammary anticancer activity and among these are melatonin and retinoic acid. NS has been examined in animals exposed to 7,12-di-methylbenz(a)anthracene (DMBA), mammary cancer causing substance which showed NS reduces the carcinogenic effects of DMBA (El-Aziz *et al.*, 2005). Inactivation of MCF-7 breast cancer cells has been demonstrated by NS extracts (Farah *et al.*, 2003).

4.1 Colon Cancer

The molecular mechanism of action of TQ in colon cancer has been suggested. Thus, colon cancer is inhibited in G1 phase cell cycle and apoptosis is mediated by TQ (Gali-Muhtasib *et al.*, 2004). The 1,2-dimethylhydrazine (DMH), colon cancer inducer, damage erythrocytes has been reported and NS has the ability to detoxify DMH (Harzallah *et al.*, 2012; Worthen *et al.*, 1998). The preneoplastic lesions for colon cancer have been investigated and found that colon cancer in post-initiation stage can be prevented by volatile components of *Nigella* seeds (Salim *et al.*, 2003).

5.0 Conclusion

The Islamic claim made by prophet Muhammad over 1400 years ago that "black seed (*Nigella sativa* Linn.) has the cure for all diseases" has a much more meaningful and acceptable dimension to it given the overwhelming scientific data obtained, as outlined in the reviews, that supports it. The alkaloids present in *Nigella sativa* Linn. could now be obtained by total synthesis and the study of their pharmacological properties should make very interesting research studies for the future.

6.0 Acknowledgements

We thank Professor K.D. Rainsford for his encouragement and helpful suggestions in writing this second review.

7.0 References

- Abdel-Fattah AM, Matsumoto K, Watanabe H. (2000) Antinociceptive effects of *Nigella sativa* oil and its major component, thymoquinone, in mice. *Eur. J Pharmacol* 400(1): 89-97.
- Aboul-El-Ela. (2002) Cytogenetic studies on *Nigella sativa* seeds extract and thymoquinone on mouse cells infected with schistosomiasis using karyotyping. *Mut Res* 516(1-2): 11-17.
- Ahmed OG, El-Mottaleb NA. (2013) Renal function and arterial blood pressure alterations after exposure to acetaminophen with a potential role of *Nigella sativa* oil in adult male rats. *J Physiol Biochem* 69(1): 1-13.

- Ahmad A, Husain A, Mujeeb M, Khan SA, Najmi AK, Siddique NA, Damanhour Z, Anwar F. (2013) A review on therapeutic potential of *Nigella sativa*: A miracle herb. Asian Pac J Trop Biomed 3(5): 337-352.
- Akhtar MS, Riffat S. (1991) Field trial of *Saussurea lappa* roots against nematodes and *Nigella sativa* seeds against cestodes in children. JPMA. J Pak Med Assoc 41(8):185-187.
- Alam S, Khan ZI, Mustafa G, Kumar M, Islam F, Bhatnagar A, Ahmad FJ. (2012) Development and evaluation of thymoquinone-encapsulated chitosan nanoparticles for nose-to-brain targeting: a pharmacoscintigraphic study. Int J Nanomed 7:5705-5718.
- Al-Ghamdi MS. (2001) The anti-inflammatory, analgesic and antipyretic activity of *Nigella sativa*. J Ethnopharm 76(1):45-48.
- Al-Ghamdi MS. (2003) Protective effect of *Nigella sativa* seeds against carbon tetrachloride-induced liver damage. Am J Chin Med 31(5):721-728.
- Al-Majed AA, Daba MH, Asiri YA, Al-Shabanah OA, Mostafa AA, El-Kashef HA. (2001) Thymoquinone-induced relaxation of guinea-pig isolated trachea. Res comm Mol Pathol Pharmacol 110(5-6):333-345.
- Al-Naggar TB, Gomez-Serranillos MP, Carretero ME, Villar AM. (2003) Neuropharmacological activity of *Nigella sativa* L. extracts. J Ethnopharm 88(1):63-68.
- Al-Suhaimi EA. (2012) Hepatoprotective and immunological functions of *Nigella sativa* seed oil against hypervitaminosis A in adult male rats. Int J Vit Nutr Res 82(4):288-297.
- Ali BH. (2004) The effect of *Nigella sativa* oil on gentamicin nephrotoxicity in rats. Am J Chin Med 32(1):49-55.
- Ali BH, Blunden G. (2003) Pharmacological and toxicological properties of *Nigella sativa*. Phytother Res 17(4):299-305.
- Al-shatwi AA. (2014) Bioactivity-guided identification to delineate the immunomodulatory effects of methanolic extract of *Nigella sativa* seed on human peripheral blood mononuclear cells. Chin J Integr Med.
- Aqel M, Shaheen R. (1996) Effects of the volatile oil of *Nigella sativa* seeds on the uterine smooth muscle of rat and guinea pig. J Ethnopharmacol 52(1):23-26.
- Arafa NM, Abdel-Rahman M, El-khadragy MF, Kassab RB. (2013) Evaluation of the possible epileptogenic activity of ciprofloxacin: the role of *Nigella sativa* on amino acids neurotransmitters. Neurochem Res 38(1):174-185.

- Badary OA. (1999) Thymoquinone attenuates ifosfamide-induced Fanconi syndrome in rats and enhances its antitumor activity in mice. *J Ethnopharmacol* 67(2):135-142.
- Badary OA, Abd-Ellah MF, El-Mahdy MA, Salama SA, Hamada FM. (2007) Anticlastogenic activity of thymoquinone against benzo(a)pyrene in mice. *Food chem toxicol* 45(1): 88-92.
- Badary OA, Abdel-Naim AB, Abdel-Wahab MH, Hamada FM. (2000) The influence of thymoquinone on doxorubicin-induced hyperlipidemic nephropathy in rats. *Toxicol* 143(3):219-226.
- Badary OA, Gamal El-Din AM. (2001) Inhibitory effects of thymoquinone against 20-methylcholanthrene-induced fibrosarcoma tumorigenesis. *Cancer Detect Preven* 25(4):362-368.
- Badary OA, Taha RA, Gamal el-Din AM, Abdel-Wahab MH. (2003) Thymoquinone is a potent superoxide anion scavenger. *Drug Chem Toxicol* 26(2):87-98.
- Badr G, Lefevre EA, Mohany M. (2011) Thymoquinone inhibits the CXCL12-induced chemotaxis of multiple myeloma cells and increases their susceptibility to Fas-mediated apoptosis. *PloS one* 6(9):e23741.
- Banerjee S, Azmi AS, Padhye S, Singh MW, Baruah JB, Philip PA, *et al.* (2010). Structure-activity studies on therapeutic potential of Thymoquinone analogs in pancreatic cancer. *Pharmaceutical research* 27(6): 1146-1158.
- Banerjee S, Kaseb AO, Wang Z, Kong D, Mohammad M, Padhye S, *et al.* (2009) Antitumor activity of gemcitabine and oxaliplatin is augmented by thymoquinone in pancreatic cancer. *Cancer Res* 69(13):5575-5583.
- Bian J, Van Wingerden M, Ready JM. (2006) Enantioselective Total Synthesis of (+)- and (-)-Nigellamine A2. *J Am Chem Soc* 128:7428-7429.
- Boskabady MH, Shirmohammadi B, Jandaghi P, Kiani S. (2004) Possible mechanism(s) for relaxant effect of aqueous and macerated extracts from *Nigella sativa* on tracheal chains of guinea pig. *BMC Pharmacol* 4:3. PMID:15070429.
- Bourgou S, Ksouri R, Bellila A, Skandrani I, Falleh H, Marzouk B. (2008) Phenolic composition and biological activities of Tunisian *Nigella sativa* L. shoots and roots. *C R Biol* 331(1):48-55.
- Burits M, Bucar F. (2000) Antioxidant activity of *Nigella sativa* essential oil. *Phytother Res* 14(5):323-328.
- Chakravarty N. (1993) Inhibition of histamine release from mast cells by nigellone. *Ann Allergy Asthma Immunol* 70(3):237-242.

- Corder C, Benghuzzi H, Tucci M, Cason Z. (2003) Delayed apoptosis upon the treatment of Hep-2 cells with black seed. *Biomed Sci Instr* **39**: 365-370.
- Daba MH, Abdel-Rahman MS. (1998) Hepatoprotective activity of thymoquinone in isolated rat hepatocytes. *Toxicol Let* **95**(1):23-29.
- de Paulis T, Hemstapat K, Chen Y, Zhang Y, Saleh S, Alagille D, et al. (2006). Substituent effects of N-(1,3-Diphenyl-1H-pyrazol-5-yl)benzamides on positive allosteric modulation of the metabotropic glutamate-5 receptor in rat cortical astrocytes. *J Med Chem* **49**:3332–3344.
- El-Daly ES. (1998) Protective effect of cysteine and vitamin E, *Crocus sativus* and *Nigella sativa* extracts on cisplatin-induced toxicity in rats. *J Pharm Belg* **53**(2):87-93.
- El Tahir KE, Ashour MM, Al-Harbi MM. (1993a) The cardiovascular actions of the volatile oil of the black seed (*Nigella sativa*) in rats: elucidation of the mechanism of action. *Gen Pharmacol* **24**(5):1123-1131.
- El Tahir KE, Ashour MM, Al-Harbi MM. (1993b) The respiratory effects of the volatile oil of the black seed (*Nigella sativa*) in guinea-pigs: elucidation of the mechanism(s) of action. *Gen Pharmacol* **24**(5):1115-1122.
- El-Abhar HS, Abdallah DM, Saleh S. (2003) Gastroprotective activity of *Nigella sativa* oil and its constituent, thymoquinone, against gastric mucosal injury induced by ischaemia/reperfusion in rats. *J Ethnopharmacol* **84**(2-3):251-258.
- El-Aziz MA, Hassan HA, Mohamed MH, Meki AR, Abdel-Ghaffar SK, Hussein MR (2005). The biochemical and morphological alterations following administration of melatonin, retinoic acid and *Nigella sativa* in mammary carcinoma: an animal model. *Int J Exp Pathol* **86**(6):383-396.
- El-Dakhakhny M, Madi NJ, Lember N, Ammon HP. (2002a) *Nigella sativa* oil, nigellone and derived thymoquinone inhibit synthesis of 5-lipoxygenase products in polymorphonuclear leukocytes from rats. *J Ethnopharmacol* **81**(2):161-164.
- El-Dakhakhny M, Mady N, Lember N, Ammon HP. (2002b) The hypoglycemic effect of *Nigella sativa* oil is mediated by extrapancreatic actions. *Planta Med* **68**(5):465-466.
- El-Dakhakhny M, Mady NI, Halim MA. (2000a) *Nigella sativa* L. oil protects against induced hepatotoxicity and improves serum lipid profile in rats. *Arzneimittel-Forschung* **50**(9): 832-836. *J Ethnopharmacol* **72**(1-2):299-304.
- Elliott EL, Bushell SM, Cavero M, Tolan B, Kelly TR. (2005) Total synthesis of nigellicine and nigelanine hydrobromide. *Org Let* **7**(12):2449-2451

- El-Saleh SC, Al-Sagair OA, Al-Khalaf MI. (2004) Thymoquinone and *Nigella sativa* oil protection against methionine-induced hyperhomocysteinemia in rats. *Inter J Cardiol* 93(1):19-23.
- Enomoto S, Asano R, Iwahori Y, Narui T, Okada Y, Singab AN, *et al.* (2001) Hematological studies on black cumin oil from the seeds of *Nigella sativa* L. *Biol Pharm Bull* 24(3):307-310.
- Farah IO, Begum RA. (2003) Effect of *Nigella sativa* (*N. sativa* L.) and oxidative stress on the survival pattern of MCF-7 breast cancer cells. *Biomed Sci Instr* 39:359-364.
- Fararh KM, Atoji Y, Shimizu Y, Shiina T, Nikami H, Takewaki T. (2004) Mechanisms of the hypoglycaemic and immunopotentiating effects of *Nigella sativa* L. oil in streptozotocin-induced diabetic hamsters. *Res Vet sci* 77(2):123-129.
- Fararh KM, Atoji Y, Shimizu Y, Takewaki T. (2002) Isulinotropic properties of *Nigella sativa* oil in Streptozotocin plus Nicotinamide diabetic hamster. *Res Vet Sci* 73(3): 279-282.
- Ganea GM, Fakayode SO, Losso JN, van nostrum CF, Sabliov CM, Warner IM. (2010). Delivery of phytochemical thymoquinone using molecular micelle modified poly(D, L lactide-co-glycolide) (PLGA) nanoparticles. *Nanotechnol* 21(28):285104
- Gali-Muhtasib H, Diab-Assaf M, Boltze C, Al-Hmaira J, Hartig R, Roessner A, *et al.* (2004). Thymoquinone extracted from black seed triggers apoptotic cell death in human colorectal cancer cells via a p53-dependent mechanism. *Int J Oncol* 25(4):857-866.
- Gheita TA, Kenawy SA. (2012) Effectiveness of *Nigella sativa* oil in the management of rheumatoid arthritis patients: a placebo controlled study. *Phytother Res* 26(8): 1246-1248.
- Gilani AH, Aziz N, Khurram IM, Chaudhary KS, Iqbal A. (2001) Bronchodilator, spasmolytic and calcium antagonist activities of *Nigella sativa* seeds (Kalonji): a traditional herbal product with multiple medicinal uses. *J Pak Med Assoc* 51(3):115-120.
- Hajhashemi V, Ghannadi A, Jafarabadi H. (2004) Black cumin seed essential oil, as a potent analgesic and antiinflammatory drug. *Phytother Res* 18(3):195-199.
- Hanafy MS, Hatem ME. (1991) Studies on the antimicrobial activity of *Nigella sativa* seed (black cumin). *J Ethnopharmacol* 34(2-3):275-278.
- Haq A, Abdullatif M, Lobo PI, Khabar KS, Sheth KV, al-Sedairy ST. (1995) *Nigella sativa*: effect on human lymphocytes and polymorphonuclear leukocyte phagocytic activity. *Immunopharmacol* 30(2):147-155.

- Haq A, Lobo PI, Al-Tufail M, Rama NR, Al-Sedairy ST. (1999) Immunomodulatory effect of *Nigella sativa* proteins fractionated by ion exchange chromatography. *Inter J Immunopharmacol* 21(4):283-295.
- Harzallah HJ, Grayaa R, Kharoubi W, Maaloul A, Hammami M, Mahjoub T. (2012) Thymoquinone, the *Nigella sativa* bioactive compound, prevents circulatory oxidative stress caused by 1,2-dimethylhydrazine in erythrocyte during colon postinitiation carcinogenesis. *Oxid Med Cell long* 2012:854065.
- Hawsawi ZA, Ali BA, Bamosa AO. (2001) Effect of *Nigella sativa* (Black Seed) and thymoquinone on blood glucose in albino rats. *Ann Saudi Med* 21(3-4):242-244.
- Hosseinzadeh H, Parvardeh S. (2004) Anticonvulsant effects of thymoquinone, the major constituent of *Nigella sativa* seeds, in mice. *Phytomed* 11(1):56-64.
- Houghton PJ, Zarka R, de las Heras B, Hoult JR. (1995) Fixed oil of *Nigella sativa* and derived thymoquinone inhibit eicosanoid generation in leukocytes and membrane lipid peroxidation. *Planta Med* 61(1):33-36.
- Iddamaldeniya SS, Wickramasinghe N, Thabrew I, Ratnatunge N, Thammitiyagodage MG. (2003) Protection against diethylnitrosoamine-induced hepatocarcinogenesis by an indigenous medicine comprised of *Nigella sativa*, *Hemidesmus indicus* and *Smilax glabra*: a preliminary study. *J Carcinogen* 2(1):6. PMID:16684351.
- Inamoto K, Katsuno M, Yoshino T, Arai Y, Hiroyaa K, Sakamotoa T. (2007) Synthesis of 3-substituted indazoles and benzoisoxazoles via Pd-catalyzed cyclization reactions: application to the synthesis of nigellicine. *Tetrahed* 63:2695–2711.
- Islam SN, Begum P, Ahsan T, Huque S, Ahsan M. (2004) Immunosuppressive and cytotoxic properties of *Nigella sativa*. *Phytother Res* 18(5):395-398.
- Kacem R, Meraihi Z. (2006) Effects of essential oil extracted from *Nigella sativa* (L.) seeds and its main components on human neutrophil elastase activity. *Yakugaku zasshi : J Pharm Soc Japan* 126(4):301-305.
- Kalus U, Pruss A, Bystron J, Jurecka M, Smekalova A, Lichius JJ, *et al.* (2003) Effect of *Nigella sativa* (black seed) on subjective feeling in patients with allergic diseases. *Phytother Res* 17(10):1209-1214.
- Kanter M, Coskun O, Korkmaz A, Oter S. (2004) Effects of *Nigella sativa* on oxidative stress and beta-cell damage in streptozotocin-induced diabetic rats. *Anatomical Record. Part A, Disc Mol Cell Evol Biol* 279(1):685-691.
- Kanter M, Meral I, Dede S, Gunduz H, Cemek M, Ozbek H, *et al.* (2003a). Effects of *Nigella sativa* L. and *Urtica dioica* L. on lipid peroxidation, antioxidant enzyme systems and some liver enzymes in CCl₄-treated rats. *J Vet Med A, Physiol Pathol Clin Med* 50(5): 264-268.

- Kanter M, Meral I, Yener Z, Ozbek H, Demir H. (2003b) Partial regeneration/proliferation of the beta-cells in the islets of Langerhans by *Nigella sativa* L. in streptozotocin-induced diabetic rats. *Tohoku J Exp Med* 201(4):213-219.
- Khader M, Bresgen N, Eckl PM. (2010) Antimutagenic effects of ethanolic extracts from selected Palestinian medicinal plants. *J Ethnopharmacol* 127(2):319-324.
- Khan MA. (1999) Chemical composition and medicinal properties of *Nigella sativa* Linn. *Inflammopharmacol* 7(1):15-35.
- Khan MA, Ashfaq MK, Zuberi HS, Mahmood MS, Gilani AH. (2003a) The in vivo antifungal activity of the aqueous extract from *Nigella sativa* seeds. *Phytother Res* 17(2):183-186.
- Khan MA, Chen HC, Tania M, Zhang DZ. (2011) Anticancer activities of *Nigella sativa* (black cumin). *African J Trad Comple Altern Med* 8(5 Suppl):226-232.
- Khan N, Sharma S, Sultana S. (2003b). *Nigella sativa* (black cumin) ameliorates potassium bromate-induced early events of carcinogenesis: diminution of oxidative stress. *Human Exp Toxicol* 22(4):193-203.
- Kirui PK, Cameron J, Benghuzzi HA, Tucci M, Patel R, Adah F, *et al.* (2004) Effects of sustained delivery of thymoquinone on bone healing of male rats. *Biomed Sci Instrument* 40:111-116.
- Korany NS, Ezzat BA. (2011) Prophylactic effect of green tea and *Nigella sativa* extracts against fenitrothion-induced toxicity in rat parotid gland. *Arch Oral Biol* 56(11):1339-1346.
- Kumara SS, Huat BT. (2001) Extraction, isolation and characterisation of antitumor principle, alpha-hederin, from the seeds of *Nigella sativa*. *Planta Med* 67(1):29-32.
- Lahm GP, Cordova D, Barry JD. (2009) New and selective ryanodine receptor activators for insect control. *Bioorg Med Chem* 17:4127-4133
- Le PM, Benhaddou-Andaloussi A, Elimadi A, Settaf A, Cherrah Y, Haddad PS. (2004) The petroleum ether extract of *Nigella sativa* exerts lipid-lowering and insulin-sensitizing actions in the rat. *J Ethnopharmacol* 94(2-3):251-259.
- Lei X, Lv X, Liu M, Yang Z, Ji M, Guo X, *et al.* (2012) Thymoquinone inhibits growth and augments 5-fluorouracil-induced apoptosis in gastric cancer cells both in vitro and in vivo. *Biochem Biophys Res Comm* 417(2):864-868.
- Mabrouk GM, Moselhy SS, Zohny SF, Ali EM, Helal TE, Amin AA, *et al.* (2002) Inhibition of methylnitrosourea (MNU) induced oxidative stress and carcinogenesis by orally administered bee honey and *Nigella* grains in Sprague Dawely rats. *J Exp clin Cancer Res* 21(3):341-346.

- Mahmood MS, Gilani AH, Khwaja A, Rashid A, Ashfaq MK. (2003) The in vitro effect of aqueous extract of *Nigella sativa* seeds on nitric oxide production. *Phytother Res* 17(8):921-924.
- Mahmoud MR, El-Abhar HS, Saleh S. (2002) The effect of *Nigella sativa* oil against the liver damage induced by *Schistosoma mansoni* infection in mice. *J Ethnopharmacol* 79(1):1-11.
- Majdalawieh AF, Hmaidan R, Carr RI. (2010) *Nigella sativa* modulates splenocyte proliferation, Th1/Th2 cytokine profile, macrophage function and NK anti-tumor activity. *J Ethnopharmacol* 131(2):268-275.
- Mansour M, Tornhamre S. (2004) Inhibition of 5-lipoxygenase and leukotriene C4 synthase in human blood cells by thymoquinone. *J Enz Inhib Med Chem* 19(5):431-436.
- Mansour MA, Ginawi OT, El-Hadiyah T, El-Khatib AS, Al-Shabanah OA, Al-Sawaf HA. (2001) Effects of volatile oil constituents of *Nigella sativa* on carbon tetrachloride-induced hepatotoxicity in mice: evidence for antioxidant effects of thymoquinone. *Res Comm Mol Pathol Pharmacol* 110(3-4):239-251.
- Mansour SW, Sangi S, Harsha S, Khaleel MA, Ibrahim AR. (2013) Sensibility of male rats fertility against olive oil, *Nigella sativa* oil and pomegranate extract. *Asian Pacific J Trop Biomed* B(7):563-568.
- Maxwell BD. (2000) The radiolabeled syntheses of JV 485, a herbicide candidate for winter wheat. *Label J Compd Radiopharm* 43:645-654
- Meral I, Kanter M. (2003) Effects of *Nigella sativa* L. and *Urtica dioica* L. on selected mineral status and hematological values in CCl4-treated rats. *Biol trace Element Res* 96(1-3):263-270.
- Meral I, Yener Z, Kahraman T, Mert N. (2001) Effect of *Nigella sativa* on glucose concentration, lipid peroxidation, anti-oxidant defence system and liver damage in experimentally-induced diabetic rabbits. *J Vet Med A, Physiol Pathol Clin Med* 48(10):593-599.
- Michelitsch A, Rittmannsberger A, Hufner A, Ruckert U, Likussar W. (2004) Determination of isopropylmethylphenols in black seed oil by differential pulse voltammetry. *Phytochem Anal* 15(5):320-324.
- Morikawa T, Xu F, Ninomiya K, Matsuda H, Yoshikawa M. (2004a) Nigellamines A3, A4, A5, and C, new dolabellane-type diterpene alkaloids, with lipid metabolism-promoting activities from the Egyptian medicinal food black cumin. *Chem Pharm Bull* 52(4):494-497.

- Morikawa T, Xu F, Kashima Y, Matsuda H, Ninomiya K, Yoshikawa M. (2004b) Novel dolabellane-type diterpene alkaloids with lipid metabolism promoting activities from the seeds of *Nigella sativa*. *Org Let* 6(6):869-872.
- Morsi NM. (2000) Antimicrobial effect of crude extracts of *Nigella sativa* on multiple antibiotics-resistant bacteria. *Acta Microbiol Pol* 49(1):63-74.
- Nair SC, Salomi MJ, Panikkar B, Panikkar KR. (1991) Modulatory effects of *Crocus sativus* and *Nigella sativa* extracts on cisplatin-induced toxicity in mice. *J Ethnopharmacol* 31(1):75-83.
- Nickavar B, Mojab F, Javidnia K, Amoli MA. (2003) Chemical composition of the fixed and volatile oils of *Nigella sativa* L. from Iran. *Z Naturforsch C* 58(9-10):629-631.
- Okuno S, Saito A, Hayashi T, Chan PH (2004). "The c-Jun N-terminal protein kinase signaling pathway mediates Bax activation and subsequent neuronal apoptosis through interaction with Bim after transient focal cerebral ischemia". *J. Neurosci.* 24 (36): 7879-87.
- Paarakh, PM. (2010) *Nigella sativa* Linn.- A comprehensive review. *Indian J Nat Prod Res* 1(4):409-429.
- Penning TD, Talley JJ, Bertenshaw SR, Carter JS, Collins WP, Docter S, et al. (1997) Synthesis and Biological Evaluation of the 1,5-Diarylpyrazole Class of Cyclooxygenase-2 Inhibitors: Identification of 4-[5-(4-Methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide (SC-58635, Celecoxib). *J Med Chem* 40:1347-1365.
- Plosker GL, Goa KL. (1991) Granisetron. A review of its pharmacological properties and therapeutic use as an antiemetic. *DRUGS* , 42(5): 805-824.
- Rastogi L, Feroz S, Pandey BN, Jagtap A, Mishra KP. (2010) Protection against radiation-induced oxidative damage by an ethanolic extract of *Nigella sativa* L. *Int J Rad Biol* 86(9):719-731.
- Ravindran J, Nair HB, Sung B, Prasad S, Tekmal RR, Aggarwal BB. (2010) Thymoquinone poly (lactide-co-glycolide) nanoparticles exhibit enhanced anti-proliferative, anti-inflammatory, and chemosensitization potential. *Biochem. Pharmacol* 79(11):1640-1647
- Rchid H, Chevassus H, Nmila R, Guiral C, Petit P, Chokairi M, et al. (2004) *Nigella sativa* seed extracts enhance glucose-induced insulin release from rat-isolated Langerhans islets. *Fundam Clin Pharmacol* 18(5):525-529.
- Saleem U, Ahmad B, Rehman K, Mahmood S, Alam M, Erum A. (2012) Nephro-protective effect of vitamin C and *Nigella sativa* oil on gentamicin associated nephrotoxicity in rabbits. *Pak J Pharm Sci* 25(4):727-730.

- Salim EI, Fukushima S. (2003) Chemopreventive potential of volatile oil from black cumin (*Nigella sativa* L.) seeds against rat colon carcinogenesis. *Nutr Cancer* 45(2): 195-202.
- Salomi MJ, Nair SC, Panikkar KR. (1991) Inhibitory effects of *Nigella sativa* and saffron (*Crocus sativus*) on chemical carcinogenesis in mice. *Nutr Cancer* 16(1):67-72.
- Salomi NJ, Nair SC, Jayawardhanan KK, Varghese CD, Panikkar KR. (1992) Antitumour principles from *Nigella sativa* seeds. *Cancer Let* 63(1):41-46.
- Samarakoon SR, Thabrew I, Galhena PB, De Silva D, Tennekoon KH. (2010) A comparison of the cytotoxic potential of standardized aqueous and ethanolic extracts of a polyherbal mixture comprised of *Nigella sativa* (seeds), *Hemidesmus indicus* (roots) and *Smilax glabra* (rhizome). *Pharmacog Res* 2(6):335-342.
- Samarakoon SR, Thabrew I, Galhena PB, Tennekoon KH. (2012) Modulation of apoptosis in human hepatocellular carcinoma (HepG2 cells) by a standardized herbal decoction of *Nigella sativa* seeds, *Hemidesmus indicus* roots and *Smilax glabra* rhizomes with anti- hepatocarcinogenic effects. *BMC Complement Alter Med* 12:25. doi:10.1186/1472-6882-12-25
- Shafi G, Munshi A, Hasan TN, Alshatwi AA, Jyothy A, Lei DK. (2009) Induction of apoptosis in HeLa cells by chloroform fraction of seed extracts of *Nigella sativa*. *Cancer Cell Int* 9: 29. doi: 10.1186/1475-2867-9-29.
- Singh A, Ahmad I, Akhter S, Jain GK, Iqbal Z, Talegaonkar S, Ahmad FJ. (2013) Nanocarrier based formulation of Thymoquinone improves oral delivery: Stability assessment, in vitro and in vivo studies *Colloids Surf., B* 102: 822–832.
- Steinmann A, Schatzle M, Agathos M, Breit R. (1997) Allergic contact dermatitis from black cumin (*Nigella sativa*) oil after topical use. *Contact dermat* 36(5):268-269.
- Swamy SM, Huat BT. (2003) Intracellular glutathione depletion and reactive oxygen species generation are important in alpha-hederin-induced apoptosis of P388 cells. *Mol Cell Biochem* 245(1-2):127-139.
- Swamy SM, Tan BK. (2000) Cytotoxic and immunopotentiating effects of ethanolic extract of *Nigella sativa* L. seeds. *J Ethnopharmacol* 70(1):1-7.
- Ulu R, Dogukan A, Tuzcu M, Gencoglu H, Ulas M, Ilhan N, *et al.* (2012) Regulation of renal organic anion and cation transporters by thymoquinone in cisplatin induced kidney injury. *Food Chem Toxicol* 50(5):1675-1679.
- Verma SK, Rastogi S, Javed K, Akhtar M, I Arorac I, Samim M. (2013) Nanothymoquinone, a novel hepatotargeted delivery system for treating CCl₄ mediated hepatotoxicity in rats. *J Mater Chem B* 1:2956–2966.

- Worthen DR, Ghosheh OA, Crooks PA. (1998) The in vitro anti-tumor activity of some crude and purified components of blackseed, *Nigella sativa* L. Anticancer Res 18(3A):1527-1532.
- Ye M, Edmunds AJF, Morris JA, Sale D, Zhanga Y, Yu J-Q. (2013) A robust protocol for Pd(II)-catalyzed C-3 arylation of (1H) indazoles and pyrazoles: total synthesis of nigellidine hydrobromide. Chem Sci 4:2374–2379.
- Yusufi M, Banerjee S, Mohammad M, Khatal S, Swamy KV, Khan EM. et al. (2013) Synthesis, characterization and anti-tumor activity of novel thymoquinone analogs against pancreatic cancer. Bioorg Med Chem Let 23:3101–3104.